

Errata for *Real Analysis for Graduate Students*,
Version 3.1

Page 16, Exercise 2.9(3): Given a set

Page 54, lines 17–24: Replace by the following.

Proposition 6.3 (1) If f is a real-valued measurable function with $0 \leq a \leq f(x) \leq b$ for all x and $\mu(X) < \infty$, then $a\mu(X) \leq \int f d\mu \leq b\mu(X)$;

(2) If f and g are measurable, real-valued, and integrable and $0 \leq f(x) \leq g(x)$ for all x , then $\int f d\mu \leq \int g d\mu$.

(3) If f is real-valued, non-negative, and integrable and c is a non-negative real number, then $\int cf d\mu = c \int f d\mu$.

(4) If $\mu(A) = 0$ and f is non-negative and measurable, then $\int f\chi_A d\mu = 0$.

Page 60, lines 7–10: Replace by the following.

Proposition 7.5 (1) If f is a real-valued measurable function with $a \leq f(x) \leq b$ for all x and $\mu(X) < \infty$, then $a\mu(X) \leq \int f d\mu \leq b\mu(X)$;

(2) If f and g are measurable, real-valued, and integrable and $f(x) \leq g(x)$ for all x , then $\int f d\mu \leq \int g d\mu$.

(3) If f is complex-valued and integrable and c is a complex number, then $\int cf d\mu = c \int f d\mu$.

(4) If $\mu(A) = 0$ and f is measurable, then $\int f\chi_A d\mu = 0$.

Proof. These follow from the definition of the Lebesgue integral of a complex-valued function, Proposition 6.3, and Theorem 7.4. For example, to prove (2), write $f = f^+ - f^-$ and $g = g^+ - g^-$. Then

$$f^+(x) - f^-(x) = f(x) \leq g(x) = g^+(x) - g^-(x)$$

implies

$$0 \leq f^+(x) + g^-(x) \leq g^+(x) + f^-(x)$$

for all x . Proposition 6.3(2) implies $\int(f^+ + g^-) \leq \int(g^+ + f^-)$, and the linearity of the Lebesgue integral implies that $\int f \leq \int g$. □

Page 63, line 1: if f and the f_n are

Page 63, Exercise 7.4, line 2: its integral is equal

Page 64, Exercise 7.14:

$$\lim_{n \rightarrow \infty} \int_0^{\infty} n e^{-nx} \frac{x^2 + 1}{x^2 + x + 1} dx.$$

Page 71, line 4: f is a Lebesgue measurable

Page 71, line 13: to $\int f$, which is finite, so taking

Page 71, line 17: A is a bounded Lebesgue measurable

Page 73, line -2: (3) Continue to suppose $\mu(X) = 1$. Deduce

Page 81, line 3: some of them. All functions in this chapter are assumed to be measurable.

Page 95, Exercise 11.7, line 4: Only one “the”

Page 97, Exercise 11.17, line 2: Suppose μ and ν are σ -finite. Prove

Page 103, line -4: (1) Note that there is at least one negative set, namely, \emptyset .

Page 119, lines 5–6: Replace “Therefore ... for some $j_0 \leq k$.” by “Let j_0 be the smallest positive integer less than or equal to k such that B_α intersects B_{j_0} .”

Page 135, line 9: $A = E_{uv} \cap (\cup_{n=1}^N (x_n - h_n, x_n))$

Page 138, line 18: Change “This implies” to “Since almost every point of $[a, c]$ is in E , this implies”

Page 145, line 4: The cases $p = 1$ and $a = 0$ are obvious, so we assume $p > 1$ and $a > 0$.

Page 146, line 21: Extraneous period at end.

Page 161, Exercise 15.30: Define $\varphi(0) = 0$ and $\varphi(x) = c_1 e^{-1/|x|^2}$ for $x \neq 0$,

Page 189, line 13: where $x \in M$

Page 196, line 2: Replace f by x twice, g by y twice